arXiv:1301.6838v1 [quant-ph] 29 Jan 2013

The mysteries of quantum correlation

Shengjun Wu

Hefei National Laboratory for Physical Sciences at Microscale and Department of Modern Physics,
University of Science and Technology of China, Hefei, Anhui 230026, China

(Dated: January 30, 2013)

Since the establishment of quantum mechanics, many fundamental problems have been stirring the hearts of physicists. Among these problems, a center issue is to understand the nature of quantum correlation. Here we reveal an essential feature of quantum correlation: its simultaneous existence in complementary bases. The classical correlation in a quantum state reaches its maximum in a certain basis and vanishes in the complementary bases, while the genuine quantum correlation does not vanish in the complementary bases. We reveal this feature of quantum correlation by defining measures based on invariance under basis change. For a bipartite quantum state, the classical correlation is the maximal correlation present in a certain optimum basis, while the quantum correlation is characterized as a series of residual correlations in mutually unbiased bases.

Quantum physics differs significantly from classical physics. A complete classical description of an object contains only information of compatible properties, while a complete quantum description of an object contains complementary information of incompatible properties simultaneously. This difference is also present in correlations. Classical correlation in a bipartite system involves correlation of only a certain property, while quantum correlation in a bipartite system involves complementary correlations of incompatible properties simultaneously (Fig.1). The simultaneous existence of complementary correlations together with the freedom in choosing which one to extract is the most important feature of quantum correlation. Schröinger introduced the word "entanglement" to describe this peculiar feature, which is termed "the spooky action at a distance" by Einstein [1–4].

Later on, entangled states are defined as the states that cannot be written as convex sums of product states. This precise definition is very helpful in terms of both mathematical and physical convenience, and it actually motivates the definition of entanglement of formation. However, this mathematical definition also conceals the essential feature of quantum correlation, for which the word "entanglement" was originally introduced. Now we know that entanglement of formation is just one particular aspect of quantum correlation. Many measures of quantum correlation have been studied, and they can be divided into two categories: one category consists of the entanglement measures [5–9], and the other consists of the measures of nonclassical correlation beyond entanglement [10–21].

Several previous entropic measures of quantum correlation are based on the difference between the quantum mutual information (which is assumed as the total correlation) and a measure of classical correlation. However, it is not clear why quantum mutual information represents the total sum of classical and quantum correlations, and why the difference between the quantum mutual information and the classical correlation represents the genuine quantum correlation. Here, we discard this assumption and reveal a very essential feature of quantum correlation. The genuine quantum correlation does not vanish under change of bases, and can be characterized as the residual correlations left in the complementary bases. Our approach is also different significantly from the approaches via Bell's inequalities [22, 23]. A Bell's inequality quantifies the complementary correlations via expectation values of complementary observables, while our approach quantifies the complementary correlations from an information-theoretical aspect.


(a)
color: brown/white
height: 5 m
weight: 300 kg
location: Hefei
velocity: 20 km/h
...
(b)
Each but only one of
$\sigma_x, \sigma_y, \sigma_z$
can be measured at a time
(c)
$\frac{1}{2}(|00\rangle\langle 00| + |11\rangle\langle 11|)$
σ_z^a perfectly correlates with σ_z^b
σ_x^a does not correlate with σ_x^b
σ_y^a does not correlate with σ_y^b
(d)
$\frac{1}{\sqrt{2}}(|01\rangle - |10\rangle)$
σ_z^a perfectly correlates with σ_z^b
σ_x^a perfectly correlates with σ_x^b
σ_y^a perfectly correlates with σ_y^b

FIG. 1. (a) A complete classical description of a classical object (a zebra) is a simple collection of information about compatible properties, such as color, height, weight, position, velocity, etc. (b) A complete quantum description (a quantum state $|\psi\rangle$) of a quantum system (a spin-1/2 particle) contains information about incompatible properties (σ_x, σ_y, σ_z) in an intrinsic way: information about incompatible properties exists simultaneously even though only a single property can be measured each time; and we can freely choose which one to measure. (c) Classical correlation in a bipartite state reaches the maximum in a certain basis, and vanishes in any complementary basis. (d) Quantum correlation in a bipartite state contains correlations in complementary bases simultaneously; and one can freely choose which basis to read out the correlation.

We start with the simplest but also most important example of entangled state for two qubits shared by Alice and Bob, known as the Einstein-Podolsky-Rosen (EPR) state or the singlet state,

$$|EPR\rangle = \frac{1}{\sqrt{2}}\left(|01\rangle - |10\rangle\right).$$

Alice can project her qubit onto the basis $\{|0\rangle, |1\rangle\}$, her measurement will leave Bob with either $|0\rangle$ or $|1\rangle$, which is perfectly anti-correlated with Alice's result. Bob can know Alice's result with certainty when he measures his qubit in the same basis, and thus achieve 1 bit (the maximum for a EPR state) of information about Alice's result. This amount is also the maximal amount of classical correlation in the EPR state. The very essential feature of the genuine quantum correlation is the simultaneous existence of correlation in complementary bases. When Alice chooses to measure her qubit in a completely different basis, Alice's measurement will leave Bob's qubit in one of two orthogonal states in the corresponding basis, and Bob can still obtain 1 bit of information about Alice's result. This fact reveals a spooky action by Alice because she can choose different bases to perform her measurement, leave Bob's qubit in different ensembles of orthogonal states, and thus Bob can always achieve 1 bit of information about Alice's result. For the EPR state, the maximum correlation is present simultaneously in all bases. We characterize the classical and the genuine quantum correlation in a rigorous way as follows.

For any bipartite quantum state ρ_{AB}, there are many measures of classical correlation [24]. Here we use the one proposed by Henderson and Vedral [25], which is also used in the definition of quantum discord [10]. Alice chooses a basis $\{|a_i\rangle_A |i = 1, \cdots, d_A\}$ of her system in a d_A-dimensional Hilbert space, and performs a measurement projecting her system onto the basis states. With probability $p_i = tr_{AB}((|a_i\rangle_A \langle a_i| \otimes I_B)\rho_{AB})$, Alice will obtain the i-th basis state $|a_i\rangle$ and Bob's system will be left in the corresponding state $\rho_i^B =_A \langle a_i| \rho_{AB} |a_i\rangle_A / p_i$. The Holevo bound of the ensemble $\{p_i; \rho_i^B\}$ that is prepared for Bob by Alice via her local measurement is given by $\chi\{\rho_{AB}|\{|a_i\rangle_A\}\} = \chi\{p_i; \rho_i^B\} \equiv S(\sum_i p_i \rho_i^B) - \sum_i p_i S(\rho_i^B)$; it denotes the upper bound of Bob's accessible information about Alice's measurement result when Alice projects her system onto the basis $\{|a_i\rangle_A\}$. The classical correlation in the state ρ_{AB} is defined as the maximal Holevo bound over all local projective measurements on Alice's system

$$C_1^h(\rho_{AB}) \equiv \max_{\{|a_i\rangle_A\}} \chi\{\rho_{AB}|\{|a_i\rangle_A\}\}. \qquad (1)$$

Let $\left\{|\mathcal{A}_i^1\rangle_A |i = 1, \cdots, d_A\right\}$ denote the optimum basis onto which Alice's system is projected so that the maximal Holevo bound is achieved.

We consider another basis $\left\{|a_j^2\rangle_A |j = 1, \cdots, d_A\right\}$ which is mutually unbiased to the basis $\left\{|\mathcal{A}_i^1\rangle_A |i = 1, \cdots, d_A\right\}$ in the sense that $|\langle a_i^1|a_j^2\rangle| = \frac{1}{\sqrt{d_A}}$, i.e., if the system is in a basis state of one basis, a projective measurement onto the mutually unbiased basis (MUB) will yield each basis state with the same probability. The most essential feature of quantum correlation is that when Alice performs a measurement in another basis $\left\{|a_j^2\rangle_A |j = 1, \cdots, d_A\right\}$ that is mutually unbiased to the optimum basis, Bob's accessible information about Alice's results, characterized by the Holevo bound, does not vanish. This residual correlation represents genuine quantum correlation, and can be used as a measure of the quantum correlation. Formally, a measure of quantum correlation $Q_2^h(\rho_{AB})$ in the state ρ_{AB} is defined as the Holevo bound of Bob's accessible information about Alice's results, maximized over all bases mutually unbiased to the basis $\left\{|\mathcal{A}_i^1\rangle_A |i = 1, \cdots, d_A\right\}$, i.e.,

$$Q_2^h(\rho_{AB}) \equiv \max_{\{|a_j^2\rangle_A\}} \chi\{\rho_{AB}|\{|a_j^2\rangle_A\}\} \qquad (2)$$

where $\left\{|a_j^2\rangle_A |j = 1, \cdots, d_A\right\}$ is any basis mutually unbiased to the optimum basis $\left\{|\mathcal{A}_i^1\rangle_A |i = 1, \cdots, d_A\right\}$ that achieves the maximal classical correlation in (1). Let $\left\{|\mathcal{A}_j^2\rangle_A |j = 1, \cdots, d_A\right\}$ denote the optimum basis that achieves the maximum quantum correlation in (2); it is mutually unbiased to the basis $\left\{|\mathcal{A}_i^1\rangle_A |i = 1, \cdots, d_A\right\}$.

Like the case in characterizing entanglement, a single quantity is not sufficient to describe the full property of quantum correlation because there could be many types of quantum correlation as well. Following the same line of reasoning, we can define the residual correlation in a third MUB as

$$Q_3^h(\rho_{AB}) \equiv \max_{\{|a_k^3\rangle_A\}} \chi\{\rho_{AB}|\{|a_k^3\rangle_A\}\} \qquad (3)$$

where $\left\{|a_k^3\rangle_A |k = 1, \cdots, d_A\right\}$ is any basis mutually unbiased to both $\left\{|\mathcal{A}_i^1\rangle_A |i = 1, \cdots, d_A\right\}$ and $\left\{|\mathcal{A}_j^2\rangle_A |j = 1, \cdots, d_A\right\}$. The optimum basis to achieve the maximum in (3) is denoted as $\left\{|\mathcal{A}_k^3\rangle_A |k = 1, \cdots, d_A\right\}$.

Suppose in this way we can define M quantities for the measures of correlation, which are conveniently written as a single correlation vector $\overrightarrow{C} \equiv (C_1^h, Q_2^h, Q_3^h, \cdots, Q_M^h)$ for the state ρ_{AB}. The number M cannot be greater than the number of MUBs that exist in the d_A-dimensional Hilbert space. The first quantity C_1^h denotes the maximal classical correlation present in the state ρ_{AB}, which can be revealed when Alice performs a measurement of her system onto the basis $\left\{|\mathcal{A}_i^1\rangle_A |i = 1, \cdots, d_A\right\}$. As classical correlation will vanish when measured in a mutually unbiased basis (see the example for CQ states), all the other quantities describe genuine quantum types of correlation. The second quantity Q_2^h denotes the maximal genuine quantum correlation, and $\left\{|\mathcal{A}_j^2\rangle_A |j = 1, \cdots, d_A\right\}$ denote the optimum basis to reveal this correlation. The third quantity Q_3^h denotes another type of genuine quantum correlation which is complementary to the first type

Q_2^h, and $\left\{\left|\mathcal{A}_k^3\right\rangle_A |j=1,\cdots,d_A\right\}$ denotes the basis to reveal the second type of quantum correlation.

Now we consider some examples to see how these measures in terms of MUBs are well justified as measures of genuine quantum correlation. For a bipartite pure state written in the Schmidt basis, $|\psi\rangle_{AB} = \sum_i \sqrt{\lambda_i}\,|a_i\rangle\,|b_i\rangle$, the maximal classical correlation can be revealed when Alice performs her measurement onto her Schmidt basis $\{|a_i\rangle\}$; thus one immediately has $C_1^h = S(\rho_B) = \sum_i -\lambda_i \log_2 \lambda_i$. If Alice chooses another basis $\{|a_i'\rangle\}$, whenever she gets a particular measurement result, Bob will be left with a pure state. Therefore, one can easily obtain the maximal true quantum correlation $Q_2^h = S(\rho_B) = C_1^h$; and any other basis will yield the same amount of quantum correlation. So the correlation vector for a bipartite pure state $|\psi\rangle_{AB}$ is given as $\overrightarrow{C} = (S(\rho_B), S(\rho_B), \cdots, S(\rho_B))$. The correlation vector shows a unique feature of the correlations in a pure state: the amount of classical correlation is equal to the amount of quantum correlation revealed in any basis, and both are equal to the von Neumann entropy of the reduced density matrix on either side, which is the usual measure of entanglement in a pure state.

A classical-quantum (CQ) state is a bipartite state that can be written as

$$\rho^{cq} = \sum_i q_i\,|i\rangle\,\langle i| \otimes \sigma_i \tag{4}$$

where $\{q_i\}$ is a probability distribution, $\{|i\rangle\,|i=0,1,\cdots,d_A-1\}$ is a basis of system A in a d_A-dimensional Hilbert space, and $\{\sigma_i\}$ is a set of density matrices of system B. The maximal classical correlation is revealed when Alice performs her measurement in the basis $\{|i\rangle\}$ [12], thus the maximal classical correlation in the CQ state ρ^{cq} is given by $C_1^h = \chi\{q_i;\sigma_i\} = S(\sum_i q_i\sigma_i) - \sum_i q_i S(\sigma_i)$. In order to calculate the amount of quantum correlation, Alice projects her system onto another basis $\left\{\left|a_j^2\right\rangle\right\}$ that is mutually unbiased to the optimum basis $\{|i\rangle\}$ for classical correlation. From $\left|\left\langle i|a_j^2\right\rangle\right|^2 = \frac{1}{d_A}$ we have

$$\begin{aligned}\left\langle a_j^2\right|\rho^{cq}\left|a_j^2\right\rangle &= \sum_i q_i \left\langle a_j^2\right|(|i\rangle\,\langle i|)\left|a_j^2\right\rangle \sigma_i \\ &= \frac{1}{d_A}\sum_i q_i\sigma_i = \frac{1}{d_A}\rho_B.\end{aligned} \tag{5}$$

For each different result j that Alice obtains, Bob is left with the same state $\rho_B = \sum_i q_i\sigma_i$; thus Bob's state has no correlation with Alice's result, and we immediately have $Q_2^h = Q_3^h = \cdots = 0$ according to the definitions of these quantities. Hence, for a CQ state, the correlation vector is given as $\overrightarrow{C} = (C_1^h, 0, \cdots, 0)$. The only correlation present in a CQ state is the classical correlation, and the quantum correlation in any MUB vanishes!

Next, we consider the Werner states of a $d\times d$ dimensional system [26],

$$\rho_w = \frac{1}{d(d-\alpha)}\,(I - \alpha P) \tag{6}$$



FIG. 2. Three measures of quantum correlation for the Werner states as functions of α when $d=2$ (left) and $d=3$ (right). The red curve represents our measure Q_2^h, the green curve represents the quantum discord Q_d and the blue curve represents the entanglement of formation E_f.

where $-1 \le \alpha \le 1$, I is the identity operator in the d^2-dimensional Hilbert space, and $P = \sum_{i,j=1}^d |i\rangle\,\langle j| \otimes |j\rangle\,\langle i|$ is the operator that exchanges A and B. Because the Werner states are invariant under unitary transformation of the form $U\otimes U$, the maximal classical correlation can be revealed when Alice simply projects her system onto the basis states $\{|i\rangle\}$. With probability $p_i = \frac{1}{d}$, Alice will get the i-th basis state $|i\rangle$, and Bob will be left with the state $\rho_i^B =_A \langle i|\,\rho_{AB}\,|i\rangle_A / p_i = \frac{1}{d-\alpha}\,(I - \alpha\,|i\rangle\,\langle i|)$. It is straightforward to have $C_1^h = \chi\left\{p_i;\rho_i^B\right\} = \log_2(\frac{d}{d-\alpha}) + \frac{1-\alpha}{d-\alpha}\log_2(1-\alpha) \equiv \chi_w$. Due to the symmetry of the Werner states, it is not difficult to show $Q_2^h = Q_3^h = \cdots = C_1^h = \chi_w$. So for the Werner state ρ_w, the correlation vector is given by $\overrightarrow{C} = (\chi_w, \chi_w, \cdots, \chi_w)$. The maximal quantum correlation in a Werner state can be revealed in any basis, and it is equal to the maximal classical correlation C_1^h. However, the correlation vector of a Werner state is different from that of a pure state because $C_1^h \le S(\rho_B) = \log_2 d$. The inequality becomes equality only when $d=2$ and $\alpha=1$, in which case the Werner state becomes a pure state $\rho_w = |EPR\rangle\,\langle EPR|$. For the Werner states, the symmetric discord is equal to the quantum discord Q_d [12] when Alice's measurement is restricted to projective measurements. The entanglement of formation E_f for the Werner states is given as $E_f(\rho_w) = h\left(\frac{1}{2}(1+\sqrt{1-[\max(0,\frac{d\alpha-1}{d-\alpha})]^2})\right)$, with $h(x) \equiv -\log_2 x - \log_2(1-x)$ [27]. The three different measures of quantum correlation, i.e., our measure of maximal quantum correlation Q_2^h, the quantum discord Q_d and the entanglement of formation, are illustrated in Fig.2 for comparison. From the figure, we know that the curve for entanglement of formation intersects the other two curves, so E_f could be larger or smaller than Q_2^h (Q_d). We also know that $Q_2^h \le Q_d$ for the Werner states. However, it is not clear whether this inequality holds for all bipartite states.

Let us consider a family of two-qubit states, where the reduced density matrices of both qubits are proportional to the identity operator. Such a state can be written in terms of Pauli matrices,

$$\rho_{AB} = \frac{1}{4}(I_2\otimes I_2 + \sum_{j,k=1}^{3} w_{jk}\sigma_j\otimes\sigma_k) \tag{7}$$



FIG. 3. Different measures of quantum correlation for two special classes of states: $\rho_1 = p\left|\psi^-\right\rangle\left\langle\psi^-\right| + (1-p)\left|\psi^+\right\rangle\left\langle\psi^+\right|$ (the left figure), and $\rho_2 = p\left|\psi^-\right\rangle\left\langle\psi^-\right| + \frac{1-p}{2}\left(\left|\psi^+\right\rangle\left\langle\psi^+\right| + \left|\phi^+\right\rangle\left\langle\phi^+\right|\right)$ (the right figure). In each figure, the red curve represents our measure Q_2^h, the green curve represents the quantum discord Q_d and the blue curve represents the entanglement of formation E_f. In the left figure, the green curve is not shown because $Q_d = Q_2^h$ for ρ_1.

where I_2 is the identity operator in the 2-dimensional Hilbert space of a qubit, w_{jk} are real numbers that satisfy certain conditions to ensure the positivity of the matrix in (7). These two-qubit states can be transformed by a local unitary transformation (that does not change the correlations) to the following form

$$\sigma_{AB} = \frac{1}{4}(I_2 \otimes I_2 + \sum_{j=1}^{3} r_j \sigma_j \otimes \sigma_j). \tag{8}$$

To ensure the positivity of the matrix in (8), the real vector $\overrightarrow{r} = (r_1, r_2, r_3)$ must lie inside or on the boundary of the regular tetrahedron that is the convex hull of the four points: $(-1,-1,-1)$, $(-1,1,1)$, $(1,-1,1)$ and $(1,1,-1)$. The singular values of the matrix w_{jk} are given by $|r_j|$. We rearrange the three numbers $\{r_1, r_2, r_3\}$ according to their absolute values, and denote the rearranged set as $\{\overline{r}_1, \overline{r}_2, \overline{r}_3\}$ so that $|\overline{r}_1| \geq |\overline{r}_2| \geq |\overline{r}_3|$. In the appendix, we show that the correlation vector is given by $\overrightarrow{C} = (\chi_1, \chi_2, \chi_3)$ where $\chi_j = 1 - h\left(\frac{1+|\overline{r}_j|}{2}\right)$ with $h(x) \equiv -\log_2 x - \log_2(1-x)$. We consider some special classes of states with only one parameter. When $r_1 = r_2 = r_3 = -\frac{\alpha}{2-\alpha}$ with $-1 \leq \alpha \leq 1$, the states in (8) become the Werner states for $d = 2$ in (6). When $r_1 = r_2 = 1 - 2p$ and $r_3 = -1$ with $0 \leq p \leq 1$, the states in (8) become $\rho_1 = p\left|\psi^-\right\rangle\left\langle\psi^-\right| + (1-p)\left|\psi^+\right\rangle\left\langle\psi^+\right|$; we obtain $C_1^h = 1$ and $Q_2^h = Q_3^h = 1 - h(p)$. When $r_1 = 1 - 2p$, $r_2 = r_3 = -p$, the states in (8) become $\rho_2 = p\left|\psi^-\right\rangle\left\langle\psi^-\right| + \frac{1-p}{2}\left(\left|\psi^+\right\rangle\left\langle\psi^+\right| + \left|\phi^+\right\rangle\left\langle\phi^+\right|\right)$; we have $C_1^h = max\{1 - h(p), 1 - h(\frac{1+p}{2})\}$, $Q_2^h = 1 - h(\frac{1+p}{2})$ and $Q_3^h = min\{1 - h(p), 1 - h(\frac{1+p}{2})\}$. Here, $|\psi^-\rangle = |EPR\rangle = \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle)$, $|\psi^+\rangle = \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle)$, and $|\phi^+\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$. Our measure of quantum correlation Q_2^h is compared with the quantum discord Q_d and the entanglement of formation E_f for ρ_1 and ρ_2 in Fig.3.

The correlation vector defined above relies on the special choice of the measure of classical correlation, it is not symmetric with respect to exchange of A and B. One may define a symmetric correlation vector, which is discussed in the appendix.

Quite different from other measures of entanglement and nonclassical correlation, our measures of quantum correlation provide a natural way to quantify the "spooky action at a distance", and reveal the mysteries of the genuine quantum correlation: the simultaneous existence of correlations in complementary bases. We hope that our results provide new insights into the understanding of quantum correlation, and quantum theory in general.

The author wishes to acknowledge support from the National Natural Science Foundation of China (Grants No. 11075148 and No. 11275181), the Chinese Academy of Sciences, and the National Fundamental Research Program.

APPENDIX

Here, we calculate the correlation vector for the family of two-qubit states given in (7). We perform the calculation in the transformed basis, with the states rewritten in (8). Without loss of generality, suppose the numbers r_j are already arranged according to $|r_1| \geq |r_2| \geq |r_3|$, then we only need to prove that $\overrightarrow{C} = (\chi_1, \chi_2, \chi_3)$ where $\chi_j = 1 - h\left(\frac{1+|r_j|}{2}\right)$. A projective measurement performed on qubit A can be written $P_\pm^A = \frac{1}{2}(I_2 \pm \overrightarrow{n} \cdot \overrightarrow{\sigma})$, parameterized by the unit vector $\overrightarrow{n}$. We have

$$p_\pm \rho_\pm^B \equiv Tr_A\left(P_\pm^A \rho_{AB}\right) = \frac{1}{2} \cdot \frac{1}{2}(I_2 \pm \sum_j n_j r_j \sigma_j). \tag{9}$$

When Alice gets ±, qubit B will be in the corresponding states $\rho_\pm^B = \frac{1}{2}\left(I_2 \pm \sum_j n_j r_j \sigma_j\right)$, each occurring with probability $\frac{1}{2}$. The entropy $S(\rho_\pm^B)$ reaches its minimum value $h\left(\frac{1+|r_1|}{2}\right)$ when $\overrightarrow{n} = (1,0,0)$. From $\rho_B = p_+\rho_+^B + p_-\rho_-^B = \frac{1}{2}I_2$ and $S(\rho_B) = 1$, we immediately have $C_1^h = 1 - h\left(\frac{1+|r_1|}{2}\right)$. The basis for Alice's projection $P_\pm^A = \frac{1}{2}(I_2 \pm \overrightarrow{n}' \cdot \overrightarrow{\sigma})$ in the definition of Q_2^h must be mutually unbiased to the basis parameterized by $\overrightarrow{n} = (1,0,0)$, therefore the unit vector $\overrightarrow{n}'$ must be in the form $\overrightarrow{n}' = (0, n_2, n_3)$. The maximum in the definition of Q_2^h is reached when $\overrightarrow{n}' = (0,1,0)$, and thus a calculation similar to that for C_1^h yields $Q_2^h = 1 - h\left(\frac{1+|r_2|}{2}\right)$. For a qubit system, three MUBs exist. We can reveal the quantum correlation in another (and the last) MUB, which corresponds to the case $\overrightarrow{n}'' = (0,0,1)$. We easily obtain $Q_3^h = 1 - h\left(\frac{1+|r_3|}{2}\right)$. For the general case when the numbers r_j are not arranged according to $|r_1| \geq |r_2| \geq |r_3|$, a similar argument yields $\overrightarrow{C} = (\chi_1, \chi_2, \chi_3)$ with $\chi_j = 1 - h\left(\frac{1+|\overline{r}_j|}{2}\right)$.

Here, we consider an alternative definition of correlation vector, which is symmetric with respect to exchange of A and B. For any bipartite quantum state ρ_{AB}, Alice chooses a basis $\{|a_i\rangle_A \,|i = 1, \cdots, d_A\}$ of her system in a d_A-dimensional Hilbert space and Bob chooses a basis
]

$\{|b_i\rangle_A\,|i=1,\cdots,d_B\}$ of his system in a d_B-dimensional Hilbert space, and each one performs a measurement projecting her/his system onto the corresponding basis states. With probability $p_{ij} = tr_{AB}((|a_i\rangle_A\langle a_i|\otimes|b_j\rangle_A\langle b_j|)\rho_{AB})$, Alice and Bob will obtain the i-th and j-th results respectively. The correlation of their measurement results are well characterized by the classical mutual information

$$I\{p_{ij}\} = H\{p_i^a\} + H\{p_j^b\} - H\{p_{ij}\} \tag{10}$$

where $H\{p_k\}$ is the Shannon entropy of the probability distribution $\{p_k\}$, and $p_i^a = \sum_j p_{ij}$, $p_j^b = \sum_i p_{ij}$ are the marginal probability distributions. The symmetric measure of classical correlation C_1^s in the state ρ_{AB} is defined as the maximal classical mutual information of the local measurement results, maximized over all local bases on both size, i.e.,

$$C_1^s(\rho_{AB}) = \max_{\{|a_i\rangle\otimes|b_j\rangle\}} I\{p_{ij}\}\,. \tag{11}$$

Let $\{|\mathcal{A}_i^1\rangle_A\,|i=1,\cdots,d_A\}$ and $\{|\mathcal{B}_i^1\rangle_B\,|i=1,\cdots,d_B\}$ denote the optimum local bases of both systems to achieve the maximal classical correlation. The symmetric measure of maximal quantum correlation Q_2^s is defined as the maximal residual correlation over all local bases that are mutually unbiased to the optimum local bases for the classical correlation C_1^s, i.e.,

$$Q_2^s(\rho_{AB}) = \max_{\{|a_i^2\rangle\otimes|b_j^2\rangle\}} I\{p'_{ij}\} \tag{12}$$

where $\{|a_i^2\rangle\}$ ($\{|b_j^2\rangle\}$) is mutually unbiased to $\{|\mathcal{A}_i^1\rangle_A\}$ ($\{|\mathcal{B}_i^1\rangle_B\}$), and $p'_{ij} = tr_{AB}((|a_i^2\rangle_A\langle a_i^2|\otimes|b_j^2\rangle_A\langle b_j^2|)\rho_{AB})$. The optimum basis to achieve $Q_2^s(\rho_{AB})$ is denoted as $\{|\mathcal{A}_i^2\rangle\otimes|\mathcal{B}_j^2\rangle\}$, which is mutually unbiased to $\{|\mathcal{A}_i^1\rangle\otimes|\mathcal{B}_j^1\rangle\}$. Similarly, Q_3^s denotes the residual quantum correlation in a third complementary basis that is mutually unbiased to both $\{|\mathcal{A}_i^1\rangle\otimes|\mathcal{B}_j^1\rangle\}$ and $\{|\mathcal{A}_i^2\rangle\otimes|\mathcal{B}_j^2\rangle\}$. In this way we have an alternative correlation vector $\overrightarrow{C_s} = (C_1^s, Q_2^s, Q_3^s, \cdots, Q_M^s)$, which is symmetric with respect to the change of A and B.

Because the Holevo bound is the upper bound of accessible classical mutual information, we immediately know from the above definitions that the asymmetric correlation vector $\overrightarrow{C}$ is an upper bound of the symmetric correlation vector $\overrightarrow{C_s}$ for each component, i.e., $C_1^h \geq C_1^s$, $Q_2^h \geq Q_2^s$, $\cdots$, $Q_M^h \geq Q_M^s$. It is not difficult to show that the asymmetric correlation vector $\overrightarrow{C}$ actually coincides with the symmetric one $\overrightarrow{C_s}$ (i.e., $\overrightarrow{C} = \overrightarrow{C_s}$) for the CQ states, the Werner states, and the two-qubit states in (7).

[1] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. **47**, 777 (1935).
[2] E. Schrödinger, Naturwissenschaften **23,** 807 (1935).
[3] E. Schrödinger and M. Born, Mathematical Proceedings of the Cambridge Philosophical Society 31 (4), 555-563 (1935).
[4] The Born-Einstein Letters: correspondence between Albert Einstein and Max and Hedwig Born from 1916 to 1955, Walker, New York, 1971.
[5] C.H. Bennett, H.J. Bernstein, S. Popescu, and B. Schumacher, Phys. Rev. A **53**, 2046 (1996).
[6] C.H. Bennett et al, Phys. Rev. A **54**, 3824 (1996).
[7] V. Vedral, et al, Phys. Rev. Lett. **78**, 2275 (1997).
[8] M. Christandl and A. Winter, J. Math. Phys. **45**, 829 (2004).
[9] R. Horodecki et al., Rev. Mod. Phys. **81**, 865-942 (2009).
[10] H. Ollivier and W. H. Zurek, Phys. Rev. Lett. **88**, 017901 (2001).
[11] M. Piani, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. **100**, 090502 (2008).
[12] S. Wu, U. V. Poulsen, and K. Mølmer, Phys. Rev. A 80, 032319 (2009).
[13] J. Oppenheim, M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. **89**, 180402 (2002).
[14] M. Horodecki, K. Horodecki, P. Horodecki, R. Horodecki, J. Oppenheim, A. Sen De, and U. Sen, Phys. Rev. Lett. **90**, 100402 (2003).
[15] S. Luo, Phys. Rev. A **77**, 022301 (2008).
[16] K. Modi, T. Paterek, W. Son, V. Vedral, and M. Williamson, Phys. Rev. Lett. **104**, 080501 (2010).
[17] B. Dakic, V. Vedral, and C. Brukner, Phys. Rev. Lett. **105**, 190502 (2010).
[18] S. Luo, and S. Fu, Phys. Rev. A. **82**, 034302 (2010).
[19] G. Adesso, and A. Datta, Phys. Rev. Lett. **105**, 030501 (2010).
[20] P. Giorda, and M.G.A. Paris, Phys. Rev. Lett. **105**, 020503 (2010).
[21] K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral, Rev. Mod. Phys. **84**, 1655-1707 (2012).
[22] J.S. Bell, Physics **1**, 195-200 (1964).
[23] M. Genovese, Phys. Rep. **413**, 319-396 (2005).
[24] S. Wu, Phys. Rev. A **86**, 012329 (2012).
[25] L. Henderson, and V. Vedral, J. Phys. A **34**, 6899 (2001).
[26] R.F. Werner, Phys. Rev. A **40**, 4277 (1989).
[27] W. K. Wootters, Quant. Inf. & Comp. **1**, 27 (2001).